



2007-2008 Pesticide Compatibility, Planting Window & Tank Mix Recommendations

ABI Extender PLUS America's Best Inoculant PLUS pesticide: <i>Maximum Planting Window Allowed After Inoculant Application</i>								
Liquid Inoculant formulations WITH ABI EXTENDER		Pesticide(s)	In Tank Slurry ¹	Maximum Planting Window (Weeks)				
Liquid Inoculant or Concentrate Liquid Inoculant	plus ABI Extender			(No Pesticide)	1 week	2 weeks	3 weeks	4 weeks
			<ul style="list-style-type: none"> • Apron-Maxx[®] RTA • Apron-Maxx[®] RFC • Warden[®] RTA 	4 hours				4 weeks
			<ul style="list-style-type: none"> • Apron-Maxx[®] Bean Pak[™] • Cruiser-Maxx[®] Pak² • SoyGard[®] L Multi-Pak • SoyGard L Multi-Pak + Gaucho^{®2} 	4 hours			3 weeks	

America's Best Inoculant PLUS pesticide: <i>Maximum Planting Window Allowed After Inoculant Application</i>											
Liquid Inoculant formulations WITHOUT Extender		Pesticide(s)	In Tank Slurry ¹	Maximum Planting Window (Days)							
Liquid Inoculant or Concentrate Liquid Inoculant				(No Pesticide)	1 day	2 days	3 days	4 days	5 days	6 days	7 days
			<ul style="list-style-type: none"> • Apron-Maxx RTA • Apron-Maxx RFC • Warden RTA • Apron-Maxx Bean Pak • SoyGard L Multi-Pak • Cruiser-Maxx Pak² • SoyGard L Multi-Pak plus Gaucho² 	4 hours					5 days		
				24 hours							7 days

America's Best Inoculant PLUS pesticide: <i>Maximum Planting Window Allowed After Inoculant Application</i>		
Dry inoculant formulations:	Inoculant only	Inoculant plus one of: Apron-Maxx RTA, Apron-Maxx RFC, Warden RTA, Apron-Maxx Bean Pak, SoyGard L Multi-Pak, Cruiser-Maxx Pak ² , or SoyGard L Multi-Pak plus Gaucho ²
Non-sterile Peat Soybean	Plant within 24 hours of application	Plant within 4 hours of application
Sterile Peat Soybean	Plant within 24 hours of application	Plant within 3 days of application

Precautions: Store treated seed in cool place, out of direct sunlight. Always use non-chlorinated water for pesticide dilutants when mixing with inoculants. Read and follow all label directions.

¹ "Slurry" refers to co-mixing liquid inoculant together with the seed treatment in the treater tank. The inoculant plus the named seed treatment should be kept together in the tank no longer than four hours before application to the seed.

² Pending label registration.

America's Best Inoculant is a registered trademark of Advanced Biological Marketing. ApronMaxx RTA, ApronMaxx RFC, Bean Pak Cruiser-Maxx are trademarks or registered trademarks Syngenta Group Company. Warden is a trademark of Agrilience LLC. SoyGard and Gaucho are registered trademarks of Gustafson LLC.